

# The Journal of **Alternative** and **Complementary Medicine**

Journal of Alternative and Complementary Medicine:  
<http://mc.manuscriptcentral.com/jaltcompmed>

## GDV Analysis of Arterial Hypertension

Journal:	<i>Journal of Alternative and Complementary Medicine</i>
Manuscript ID:	JACM-2010-0084
Manuscript Type:	Original Articles
Date Submitted by the Author:	10-Feb-2010
Complete List of Authors:	Yakovleva, Ekaterina; Russian Federal Medical University, Medical cybernetics Zarubina, Tatiana; Russian Federal Medical University, Medical cybernetics Strychkov, Peter; Institute of Advanced Education, Functional diagnostic Kovelkova, Margarita; Russian Federal Medical University, Medical cybernetics Aleksandrova, Elena; Institute of Advanced Education, Functional diagnostic
Keywords:	biofield, bioenergy
Abstract:	<p>Objectives: To study potentials of Electrophotonic - Gas Discharge Visualization (EPC/GDV) analysis for detecting patients with arterial hypertension of different degree of severity in the course of population screening.</p> <p>Design: To assess dependence of the most qualitative patient grouping on one of the recognized arterial hypertension (AH) classifications, discriminative functions for detecting patients with different degrees and stages of AH as well as the risk of cardio-vascular complications were calculated. Dependence of discriminative functions on patients' gender was studied. The model of logistic regression for detecting patients with different degrees of arterial hypertension was built.</p> <p>Procedures: Participants were 603 patients, aged from 18 to 83, 265 males and 338 females. All participants were classified into groups according to degree and stage of arterial hypertension and possible risk of cardio-vascular complications in the nearest 10 years. In the course of the study the following GDV parameters were analyzed: image area, normalized area, intensity, spectrum width, brightness and fractality.</p> <p>Results: The study resulted in calculation of discriminative functions for detecting patients with various degrees, stages of arterial hypertension and risk of cardio-vascular complications. The number</p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

	<p>of diagnostic parameters increased in accordance with the degree of AH manifestations from 8 to 22, and specificity and sensitivity of the obtained functions made up from 70% to 80%. Classification of patients according to gender increased diagnostic accuracy by 5-9%.</p> <p>Conclusions: Reliable differences between the control group (healthy patients) and groups with various degrees and stages of AH were calculated with sufficiently high accuracy which allowed to include Electrophotonic - Gas Discharge Visualization technique into the population screening.</p>



For Peer Review

## GDV Analysis of Arterial Hypertension

ELENA V. ALEKSANDROVA<sup>1</sup>, TATIANA V. ZARUBINA<sup>2</sup>, MARGARITA N. KOVELKOVA<sup>2</sup>, PETER V.STRYCHKOV<sup>1</sup>,  
EKATERINA G. YAKOVLEVA<sup>2</sup>

1 - Institute of Advanced Education, Moscow, Russia; 2 – Russian Federal Medical University, Moscow;

E-mail: k\_iakov@mail.ru

### ABSTRACT

**Objectives:** To study potentials of Electrophotonic - Gas Discharge Visualization (EPC/GDV) analysis for detecting patients with arterial hypertension of different degree of severity in the course of population screening.

**Design:** To assess dependence of the most qualitative patient grouping on one of the recognized arterial hypertension (AH) classifications, discriminative functions for detecting patients with different degrees and stages of AH as well as the risk of cardio-vascular complications were calculated. Dependence of discriminative functions on patients' gender was studied. The model of logistic regression for detecting patients with different degrees of arterial hypertension was built.

**Procedures:** Participants were 603 patients, aged from 18 to 83, 265 males and 338 females. All participants were classified into groups according to degree and stage of arterial hypertension and possible risk of cardio-vascular complications in the nearest 10 years. In the course of the study the following GDV parameters were analyzed: image area, normalized area, intensity, spectrum width, brightness and fractality.

**Results:** The study resulted in calculation of discriminative functions for detecting patients with various degrees, stages of arterial hypertension and risk of cardio-vascular complications. The number of diagnostic parameters increased in accordance with the degree of AH manifestations from 8 to 22, and specificity and sensitivity of the obtained functions made up from 70% to 80%. Classification of patients according to gender increased diagnostic accuracy by 5-9%.

**Conclusions:** Reliable differences between the control group (healthy patients) and groups with various degrees and stages of AH were calculated with sufficiently high accuracy which allowed to include Electrophotonic - Gas Discharge Visualization technique into the population screening.

### INTRODUCTION

Arterial hypertension (AH) is the most common chronic disease in the world and one of the most serious medico-social problems. 30% of adult population of industrial countries present with elevated degree of arterial pressure and only 12-15% of population having stable arterial hypertension. The disease is a common cause of life-endangering acute heart (myocardial infarction) and brain (cerebral stroke) diseases including people of the working age. About 50% of all fatal cases from cardio-vascular diseases fall on arterial hypertension. It is the underlying cause of numerous chronic diseases of the heart, brain, kidneys, eyes and other so-called target organs.

Even today, when doctors have sufficient number of potent antihypertensive at their disposal average life span of the patients with arterial hypertension is not high. Arterial pressure often elevates among adolescents, the disease dramatically getting younger along with most cardiovascular diseases. Hypertension and atherosclerosis are turning into important cause of premature death<sup>1</sup>.

There is no doubt that prophylaxis and treatment of arterial hypertension are most effective at its early stages and not at the chronic stage with organic changes. It is disappointing that in many cases the onset of the disease stays unseen as early rises in arterial pressure (AP) are not always associated with subjective symptoms which makes arterial hypertension a disease difficult to diagnose at the early stages. So the early diagnostics of arterial hypertension and a special program of its further development prevention present an important scientific and practical problem<sup>2</sup>. The focus today is on methods allowing detecting early changes characteristic for arterial hypertension. Public health service needs highly sensitive non-invasive screening tests for AH.

1  
2  
3 One of the promising techniques of evaluating the patient's condition is Electrophotonic - Gas  
4 Discharge Visualization (EPC/GDV) i.e. computerized registration and analysis of photoelectronic  
5 emission stimulated by electromagnetic field and intensified in gas discharge. Glow of different objects  
6 in electromagnetic fields was discovered over 200 years ago, but only development of the GDV  
7 instruments in 1995 by Dr K. Korotkov team opened-up perspectives of medical applications for this  
8 approach. Today this technique acquires wider and wider use for diagnostics of different diseases<sup>3-10</sup>.  
9 Significant results obtained in detecting AH with GDV technique<sup>10-14</sup> demonstrate perspective of  
10 further development in this field.

11 One of the major current theories on the etiology of hypertension still remains neurogenic theory<sup>1-2</sup>.  
12 As psychoemotional strain is mainly realized through intense production and liberation of  
13 catecholamines, another concept – of raising the tone of sympathetic nervous system as key  
14 mechanism of AH development – is believed to be significant as well. Indeed, findings of  
15 cardiointervalography prove higher tone of sympathetic nervous system in most hypertensive patients.  
16 The authors of GDV technique, on the basis of conducted investigations, believe that GDV-grams reflect  
17 the activity of vegetative nervous system (V.N.S.), the balance of its sympathetic and parasympathetic  
18 divisions<sup>3-4</sup>. That allows to suggest that GDV-gram evaluation is likely to detect patients with arterial  
19 hypertension.

20 Thus, the objective of the given study was evaluation of GDV potential for detecting patients with  
21 different degrees of severity of arterial hypertension in the course of population screening.

22 Several classifications of arterial hypertension are accepted in medicine. To date, according to UHO  
23 recommendations arterial hypertension is classified into 3 degrees (depending on the degree of AP  
24 elevation) and 3 stages (depending on the involvement of target organs). There are also 4 categories of  
25 arterial hypertension depending on the likelihood of cardio-vascular complications in the nearest 10  
26 years. The complications are related to the presence of the risk-factors, involvement of target organs  
27 and/or concomitant (associated) diseases<sup>2</sup>.

28 The study was aimed at:

- 29 1. Calculating discriminative functions to detect patients with different degrees and stages of  
30 arterial hypertension as well as the risk of cardio-vascular complications; assessing  
31 dependence of the most qualitative patient grouping on one of the recognized AH  
32 classifications.
- 33 2. Assessing the influence of patients' gender on calculation discriminative functions.
- 34 3. Building the model of logistic regression to detect patients with different degrees of AH  
35 severity.

### 36 **Materials and methods**

37 603 patients aged from 18 to 83, 265 males and 338 females served as participants. All were divided  
38 into groups according to AH degree and stage and degree of cardio-vascular complications risk in the  
39 nearest 10 years. Groups were divided as follows:

40 Control group – 136 people (47 men and 89 women); diagnosed AH1 of the 1<sup>st</sup> degree – 92 persons  
41 (38 men and 54 women); diagnosed AH2 of the 2<sup>nd</sup> degree – 185 people (89 men and 96 women);  
42 diagnosed AH3 of the 3<sup>d</sup> degree – 190 people (91 men and 99 women). 4 groups were also selected  
43 according to AH stage: control group – 136 people (47 men and 89 women); diagnosed AHs1 of the 1<sup>st</sup>  
44 stage – 103 people (40 men and 63 women); diagnosed AHs2 of the 2<sup>nd</sup> stage – 283 people (130 men  
45 and 153 women); diagnosed AHs3 of the 3<sup>d</sup> stage – 81 persons (48 men and 33 women).

46 Another 5 groups were selected according to the likely risk of cardio-vascular complications: control  
47 group – 136 people (47 men and 89 women); low risk (risk 1) – 56 people (24 men and 32 women);  
48 moderate risk (risk 2) – 88 people (33 men and 55 women); high risk (risk 3) – 114 people (51 men and  
49 63 women); extremely high risk (risk 4) – 209 people (110 men and 99 women).

50 In the course of the study the following EPC/GDV-gram parameters were analyzed: image area,  
51 normalized area, intensity, spectrum width, brightness and fractality. According to these parameters we  
52 analyzed images of all 10 fingers as a whole as well as of separate sectors selected in accordance with  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Korotkov's Diagnostic Table<sup>3</sup>: cerebral cortex, cerebral vessels, the right and left heart, vascular system, coronary vessels, hypophysis, hypothalamus, epiphysis, thyroid, suprarenals, kidneys, the nervous system. Organs and systems of organs which were involved in the onset and progress of arterial hypertension were under consideration.

All data were processed with the "GDV-Processor" program to calculate abovementioned parameters; discriminative functions were calculated with the help of step by step discriminative analysis in «SPSS Statistics 17.0» and «Statistica 6.0» programs.

While using different diagnostic methods, one should know how reliable it is, taking into account its specificity and sensitivity for practical evaluation of different health problems. Specificity implies the share of healthy people found healthy in the course of diagnostics from the total number of healthy. Sensitivity implies the share of ill patients found ill in the course of diagnostics from the total number of ill patients.

### Results and discussion

The first stage of the work included step by step discriminative analysis including the control group and each of the three groups of arterial hypertension (according to degree of severity) separately. Results are presented in Table 1. The given figures are the result of cross-testing. The latter implies that each test is classified according to functions obtained in all test but the particular one.

As an example we present discriminative function between the control group and the group of the 1<sup>st</sup> degree of AH severity. 8 parameters were included into the obtained discriminative function, among them were spectrum width of the images of the right thumb, sector of the head (cortex and vessels), suprarenals, thyroid and kidneys.

From stage to stage of arterial hypertension the number of diagnostic parameters increases from 8 up to 19, which is understandable as it coincides with the degree of involvement of target-organs. All group proved to have the following sectors in common: cerebral cortex, thyroid and kidneys and starting with Group 2 (diagnosed AH2 of the 2<sup>nd</sup> degree) – the heart sector. Specificity and sensitivity increased along with higher degree of arterial hypertension.

It is known that GDV parameters are dependent of the patient's gender and arterial hypertension takes a different course in males and females<sup>2-5</sup>. Discriminative functions were calculated for all three degrees of arterial hypertension for males and females separately (Table 2).

Common diagnostic parameters were found for all groups, they included sectors of the cerebral cortex, vascular system, heart, thyroid and kidneys. Specificity and sensitivity of the functions obtained in groups divided on the basis of patients' gender were 5-9 % higher than for the mixed group.

The number of diagnostic parameters used and percentage of difference for females with AH1 and AH2 were found higher than that for males. Men are known to have a tendency for higher arterial pressure against women of reproductive age. Differences on AP between men and women disappear after women's menopause or ovariectomy. AH incidence is lower in women below 60 and higher over 60 against men of similar age.

The next stage of the investigation was comparison of data obtained for patients of the control group and groups with different stages of arterial hypertension.

Specificity and sensitivity values of calculated discriminative functions increased from the first stage of arterial hypertension to the third. Increase in specificity from stage to stage amounted to 67-80 %, and sensitivity – to 70-77%. The number of diagnostic parameters increased from stage 1 to stage 3 from 7 to 22. For all stages sectors of the cerebral cortex, heart, suprarenals and thyroid were involved in calculation.

Discriminative functions were also calculated for men and women separately (Table 4).

For this classification mean percentage of correct placements for men exceeded the one for women by 6-8%. It is likely to be associated with specific involvement of target organs for men and women.

Arterial hypertension is one of the main risk factors of cardiovascular diseases in women. Though the level of arterial pressure for women of the premenopause period is lower than for men of the corresponding age, AH incidence for elderly women is higher.

To sum it up, the course of arterial hypertension for women of the postmenopause period has the following characteristics:

- the disease is frequently manifested in perimenopause period (from the onset of initial climacteric symptoms to one year after the last menstruation);
- higher sensitivity to table salt;
- low-renin forms of arterial hypertension;
- narrower diameter of the aorta;
- characteristic rise in systolic arterial pressure;
- high rate of heart contractions;
- menopause associated with increase in cardiovascular response induced by stress and rise in arterial pressure registered during 24-hour monitoring of arterial pressure;
- more frequent involvement of target organs;
- more frequent hypertrophy of the left ventricle mostly of the concentric type;
- more frequent complications.

Elevated arterial pressure and higher incidence of arterial hypertension in postmenopause are believed to result from other factors as well e.g. increase in the volume of circulating blood, higher body mass and higher noradrenalin level in the blood.

Discriminative functions were calculated to detect patients with different risk of cardio-vascular complications after arterial hypertension. Specificity and sensitivity of calculated functions amounted to 64 – 73.5% and 62.2 – 76% respectively which correlates with values for the groups classified on the basis of AH degree of severity. The number of diagnostic parameters increased depending on the risk group. Sectors reflecting the heart and kidneys were found as the most common for all groups (Tables 5,6).

Results of discriminative analysis are presented on graphs Fig.1-3.

Feasibility of detecting patients with different AH degrees was tested with the help of logistic regression. Specificity and sensitivity were found close in values to those obtained by discriminative analysis but values of specificity calculated by logistic regression were somewhat lower than those calculated by discriminative analysis which determined our choice of the latter in our investigations (Table 7).

Our findings correlate well with those obtained earlier by other investigators with the help of neuron network method<sup>15</sup> as well as during comparison of the diagnosis made with the help of GDV and other diagnostic methods widely used in modern medicine<sup>16</sup>.

### Conclusions

Discriminative functions were calculated to detect patients with various degrees, stages and risk of cardio-vascular complications in case of arterial hypertension. The number of diagnostic parameters



1  
2 increased with higher degree of AH manifestation from 8 to 22 while specificity and sensitivity of  
3 calculated functions made up about 70%  
4

5  
6 Sectors of cerebral cortex, heart, thyroid, suprarenals and kidneys proved to be the most frequent  
7 diagnostic parameters. We believe that patients' grouping was most qualitative under classification  
8 according to AH stages, which may be explained by the fact that in grouping according to AH stages both  
9 degree of AP elevation and involvement of target organs were taken into account. Classification  
10 according to patients' gender increased the accuracy of diagnostics by 5-9% which was due to  
11 differences in development and course of arterial hypertension in women and men.  
12  
13

14  
15 Thus, reliable differences between the control group of healthy patients and groups with various AH  
16 degrees and stages were calculated with sufficiently high degree of accuracy which allows to include  
17 Electrophotonic - Gas Discharge Visualization technique into the population screening.  
18  
19

### 20 21 Disclosure Statement

22 The authors state that no competing financial interests exist.  
23  
24

### 25 26 References

- 27  
28 1. Makolkin V. I., Podzolkov V. I. Hypertension, Moscow., Publishing House "Medicine", 2000 (in  
29 Russian).  
30 2. Guidelines on hypertension/ Edited by Chazov Y.I., Moscow., Publishing House "Medicine",  
31 2005 (in Russian).  
32 3. Korotkov K. Human Energy Field: study with GDV bioelectrography. Backbone publishing, NY.  
33 2002.  
34 4. Measuring Energy Fields: State of the Art. GDV Bioelectrography series. Vol. I. Korotkov K. (Ed.).  
35 Backbone Publishing Co. Fair Lawn, USA, 2004. 270 p  
36 5. Polushin J, Levshankov A, Shirokov D, Korotkov K. Monitoring Energy Levels during treatment  
37 with GDV Technique. J of Science of Healing Outcome.. 2:5. 5-15, 2009  
38 6. Rgeusskaja G.V., Listopadov U.I. Medical Technology of Electrophotonics – Gas Discharge  
39 Visualization - in Evaluation of Cognitive Functions. J of Science of Healing Outcome. V.2, N 5,  
40 pp.15-17, 2009  
41 7. Rubik B., Brooks A. Digital High-Voltage Electrophotographic Measures of the Fingertips of  
42 Subjects Pre and Post-Qigong. Evidence Based Integrative Medicine. 2 (4), 245-242, 2005.  
43 8. Korotkov K., Williams B., Wisneski L. Biophysical Energy Transfer Mechanisms in Living Systems:  
44 The Basis of Life Processes. J of Alternative and Complementary Medicine, 2004, 10, 1, 49-57.  
45 9. Russo M. Russo M, Choudhri AF, Whitworth G, Weinberg AD, Bickel W, and Oz MC. Quantitative  
46 analysis of reproducible changes in high-voltage electrophotography. J of Alternative and  
47 Complementary Medicine 2001, 7, 6, 617-629.  
48 10. Measuring the Human Energy Field: State of the Science. Ed. R.A. Chez. National Institute of  
49 Health, Samueli Institute, Maryland, 2002.  
50 11. Alexandrova R, Fedoseev G, Korotkov K, Philippova N, Zayzev S, Magidov M, Petrovsky I. Analysis  
51 of the bioelectrograms of bronchial asthma patients. Proceedings of the St. Petersburg Federal  
52 Medical University; 2001;8,1:73-78.  
53 12. Yakovleva E G, Struchkov P V, Zarubina T. V. Application of Gas Discharge Visualization far early  
54 detection of patients with arterial hypertension in the course of regular check-ups.// XII all-  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Russia congress of cardio-vascular surgeons, November 25-28, 2007, Moscow. Bulletin of RCCVS  
4 named after Bakulev RAMS, v 8, n 6 november – december 2007, p 241.
- 5 13. Yakovleva E G, Struchkov P V, Kondratova E. Yu. GDV Diagnostic Potential for detecting patients  
6 with major cranial arterial hypertension and left ventricular hypertrophy / World Forum  
7 “Information technologies and Society” – 2008, - Kemer, Turkey. M. 2008, pp 164-165.
- 8 14. Yakovleva E G, Struchkov P V, Zarubina T. V., Kovelkova M. N., Alexandrova E. V. Different  
9 statistic approaches for processing results of Gas Discharge Visualization. In: XIII International  
10 Scientific Congress “Science. Information. Spirit” July 4-5, 2009, St. Petersburg, pp 56-57.
- 11 15. Volkov A., Telesheva T., Gursky V., Krighanovsky S. Statistical model of the patient’s diagnosis on  
12 the basis of his GDV-gram parameters. In: Proceedings of IX International Scientific Congress on  
13 Bioelectrography. “Science. Information. Spirit” – St. Petersburg. 2005. p 5.
- 14 16. Korotkov K., Matravers P., Pharm D., Orlov D. Review of publications on the use of Gas  
15 Discharge Visualization (GDV) technique in medicine. In: Proceedings of XII International  
16 Scientific Congress on Bioelectrography. “Science. Information. Spirit” – St. Petersburg. 2008. pp  
17 6-8.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Table 1. Grouping according to degree of AH. Results for all patients.

AH degree	specificity	sensitivity
AH1	67,6 %	62,0%
AH2	68,4%	66,5%
AH3	72,8%	77,9%

Table 2. Grouping according to AH degree. Results for males and females.

AH degree	Females		Males	
	Specificity	Sensitivity	Specificity	Sensitivity
AH1	76,4%	77,8%	80,9%	73,7%
AH2	74,2%	81,3%	63,8%	70,8%
AH3	75,3%	74,7%	78,7%	80,2%

Table 3. Grouping according to AH stages. Results for the whole group.

AH stage	Specificity	Sensitivity
Ahs1	66,9 %	70,9%
Ahs2	67,6%	73,5%
AHs3	80,1%	76,5%

Table 4. Grouping according to AH stages. Results for males and females.

AH stage	females		males	
	Specificity	Sensitivity	Specificity	Sensitivity
Ahs1	80,9%	64,1%	80,9%	79,5%
Ahs2	68,1%	81,3%	75,3%	75,8%
AHs3	72,3%	67,3%	83,1%	84,4%

Table 5. Grouping according to the risk of cardio-vascular complications. Results for the mixed group.

Risk of cardio-vascular complications	Specificity	Sensitivity
Risk 1	73,5 %	62,5%
Risk 2	69,9%	72,7%
Risk 3	64,0%	64,0%
Risk 4	69,9%	76,1%

Table 6. Grouping according to the risk of cardio-vascular complications. Results for males and females.


Risk of cardio-vascular complications	females		males	
	Specificity	Sensitivity	Specificity	Sensitivity
Risk 1	82,0%	75,0%	76,6%	58,3%
Risk 2	78,7%	74,5%	70,2%	66,7%
Risk 3	74,2%	77,8%	74,5%	78,4%
Risk 4	76,4%	75,8%	68,1%	72,7%

Table 7. Grouping according to AH degrees. Results calculated by logistic regression.

AH degree	specificity	sensitivity
AH1	79,4 %	70,6%
AH2	68,3%	61,0%
AH3	58,1%	83,7%

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


Fig.1. Grouping according to degree of AH.



254x190mm (96 x 96 DPI)


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Fig.2. Grouping according to the stage of AH.



254x190mm (96 x 96 DPI)

Fig.3. Grouping according to the risk of cardio-vascular complications.



254x190mm (96 x 96 DPI)